Arkady Khoutorsky<sup>a,b,c,d,1</sup>, Robert E. Sorge<sup>d,e,1</sup>, Masha Prager-Khoutorsky<sup>f,1</sup>, Sophie Anne Pawlowski<sup>d,g</sup>, Geraldine Longo<sup>d,g</sup>, Seyed Mehdi Jafarnejad<sup>a,b</sup>, Soroush Tahmasebi<sup>a,b</sup>, Loren J. Martin<sup>d,e</sup>, Mark H. Pitcher<sup>h</sup>, Christos G. Gkogkas<sup>i</sup>, Reza Sharif-Naeini<sup>d,j</sup>, Alfredo Ribeiro-da-Silva<sup>d,g</sup>, Charles W. Bourque<sup>f</sup>, Fernando Cervero<sup>c,d</sup>, Jeffrey S. Mogil<sup>d,e,2</sup>, and Nahum Sonenberg<sup>a,b,2</sup>

<sup>a</sup>Department of Biochemistry, McGill University, Montreal, QC, Canada H3A 1A3; <sup>b</sup>Rosalind and Morris Goodman Cancer Research Centre, McGill University, Montreal, QC, Canada H3A 1A3; <sup>c</sup>Department of Anesthesia, McGill University, Montreal, QC, Canada H3G 1Y6; <sup>d</sup>Alan Edwards Centre for Research on Pain, McGill University, Montreal, QC, Canada H3G 1Y6; <sup>e</sup>Department of Psychology, McGill University, Montreal, QC, Canada H3A 1B1; <sup>f</sup>Center for Research in Neuroscience, McGill University, Montreal General Hospital, Montreal, QC, Canada H3G 1A4; <sup>g</sup>Department of Pharmacology and Therapeutics, McGill University, Montreal, QC, Canada H3G 1Y6; <sup>h</sup>National Center for Complementary and Alternative Medicine, National Institutes of Health, Porter Neuroscience Research Center, MD 20892; <sup>i</sup>Patrick Wild Centre and Centre for Integrative Physiology, University of Edinburgh, Edinburgh EH8 9XD, United Kingdom; and <sup>j</sup>Department of Physiology, McGill University, Montreal, QC, Canada H3G 1Y6

Contributed by Nahum Sonenberg, August 25, 2016 (sent for review November 19, 2015; reviewed by Eric Klann and Mark J. Zylka)

A response to environmental stress is critical to alleviate cellular injury and maintain cellular homeostasis. Eukaryotic initiation factor 2 (eIF2) is a key integrator of cellular stress responses and an important regulator of mRNA translation. Diverse stress signals lead to the phosphorylation of the  $\alpha$  subunit of eIF2 (Ser51), resulting in inhibition of global protein synthesis while promoting expression of proteins that mediate cell adaptation to stress. Here we report that  $elF2\alpha$  is instrumental in the control of noxious heat sensation. Mice with decreased eIF2 $\alpha$  phosphorylation (eIF2 $\alpha^{+/S51A}$ ) exhibit reduced responses to noxious heat. Pharmacological attenuation of eIF2a phosphorylation decreases thermal, but not mechanical, pain sensitivity, whereas increasing  $elF2\alpha$  phosphorylation has the opposite effect on thermal nociception. The impact of  $elF2\alpha$ phosphorylation (p-eIF2α) on thermal thresholds is dependent on the transient receptor potential vanilloid 1. Moreover, we show that induction of eIF2 $\alpha$  phosphorylation in primary sensory neurons in a chronic inflammation pain model contributes to thermal hypersensitivity. Our results demonstrate that the cellular stress response pathway, mediated via p-eIF2α, represents a mechanism that could be used to alleviate pathological heat sensation.

pain | eIF2a | cellular stress response pathway | TRPV1

Downloaded from https://www.pnas.org by UNIVERSITY OF TORONTO on July 2, 2022 from IP address 142.150.190.39.

**R**esponse to stress is a major cellular function involved in many physiological and pathological conditions. Cells respond to various forms of stress by activating specific molecular cascades that orchestrate antistress responses or induce apoptosis (1). A key effector of cellular stress responses is the eukaryotic initiation factor 2 (eIF2) (2). Phosphorylation of eIF2 causes a reduction in global translation, allowing cells to conserve energy and modify gene expression to effectively manage stress conditions. Diverse stress signals converge onto eIF2 to integrate stress responses through phosphorylation of the  $\alpha$  subunit of eIF2.

eIF2 binds GTP and the initiator methionyl-tRNA (Met-tRNA<sub>i</sub>) to form the ternary complex (eIF2-GTP-Met-tRNA<sub>i</sub>). The ternary complex binds the small ribosomal subunit to form the ribosomal preinitiation complex, which scans the 5'UTR of the mRNA for the start codon to initiate mRNA translation (3). On engagement of the initiation codon, GTP is hydrolyzed to GDP (4). The recycling of inactive GDP-bound eIF2 to the active GTP-bound form is catalyzed by the guanine nucleotide exchange factor, eIF2B. Phosphorylation of the  $\alpha$  subunit of eIF2 at serine 51 converts eIF2 from a substrate to a competitive inhibitor of eIF2B (4). Because the amount of eIF2B is lower than eIF2, phosphorylation of a small fraction of the eIF2 in the cell is sufficient to strongly inhibit eIF2B activity and translation initiation.

eIF2 $\alpha$  is phosphorylated by four eIF2 $\alpha$  kinases, each activated in a different stress condition (5–7). PKR (double-stranded RNA-dependent protein kinase) is activated by double-stranded RNA during viral infection; PERK (PKR-like ER kinase) by endoplasmic reticulum stress; GCN2 (general control non– derepressible-2) by nutrient deprivation and UV light; and HRI (heme-regulated inhibitor) by heme deficiency. The eIF2 $\alpha$  kinases, except for HRI, are prominently expressed in the mammalian nervous system (8).

Phosphorylation of eIF2a blocks general translation but paradoxically stimulates translation of mRNAs that contain upstream ORFs (uORFs) in their 5' UTR, such as ATF4 [a cAMP-response element binding protein 2 (CREB-2)] and CHOP (a proapoptotic transcription factor) (9). ATF4 enhances the expression of a related transcription factor, ATF3, which together with ATF4 contribute to stress adaptation by regulating genes involved in metabolism, the cellular redox status, and apoptosis (10, 11). In neurons, an activitydependent decrease in eIF2 $\alpha$  phosphorylation augments long-term potentiation (LTP) and memory via suppression of ATF4 expression (12). Conversely, up-regulation of p-eIF2 $\alpha$  is associated with long-term depression (LTD) (13, 14) and several pathophysiological conditions including viral infection, inflammation, and neurodegeneration (15–18). Elevated phosphorylation of eIF2 $\alpha$  has been documented in the brain of aged animals (19) and Alzheimer's disease patients and model mice (20, 21). Normalization of p-eIF2 $\alpha$ in Alzheimer's disease model mice rescued deficits in protein synthesis, synaptic plasticity, and spatial memory (22). Additionally,

## Significance

Distinct cellular stresses converge on the translation initiation factor, eukaryotic initiation factor  $2\alpha$  (eIF2 $\alpha$ ) to modulate the rate of protein synthesis. Increased phosphorylation of eIF2 $\alpha$ has been described in peripheral neurons from neuropathic and diabetic rats. However, the role of eIF2 $\alpha$  phosphorylation in pain has not been reported. Here we show that phosphorylation of eIF2 $\alpha$  controls thermal, but not mechanical, sensation via modulation of the activity of a major heat transducer, transient receptor potential vanilloid 1. We also find that chronic inflammation-induced thermal hypersensitivity. These results demonstrate that eIF2 $\alpha$  phosphorylation plays a major role in controlling noxious heat sensitivity.

Author contributions: A.K., R.E.S., S.A.P., G.L., S.M.J., L.J.M., M.H.P., C.G.G., R.S.-N., A.R.-d.-S., C.W.B., F.C., J.S.M., and N.S. designed research; A.K., R.E.S., M.P.-K., S.A.P., G.L., S.T., L.J.M., and M.H.P. performed research; A.K., R.E.S., M.P.-K., S.A.P., and G.L. analyzed data; and A.K., R.E.S., M.P.-K., S.A.P., S.M.J., S.T., C.G.G., R.S.-N., A.R.-d.-S., C.W.B., J.S.M., and N.S. wrote the paper.

Reviewers: E.K., New York University; and M.J.Z., University of North Carolina at Chapel Hill.

The authors declare no conflict of interest.

<sup>&</sup>lt;sup>1</sup>A.K., R.E.S., and M.P.-K. contributed equally to this work.

<sup>&</sup>lt;sup>2</sup>To whom correspondence may be addressed. Email: jeffrey.mogil@mcgill.ca or nahum. sonenberg@mcgill.ca.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10. 1073/pnas.1614047113/-/DCSupplemental.

phosphorylation of  $eIF2\alpha$  is associated with synaptic deficits and neuronal loss in prion-disease model mice (23).

The role of eIF2 in the pain pathway is unknown. An endoplasmic reticulum (ER) stress response is induced in the peripheral nervous system of type I diabetic rats, as phosphorylation of PERK and eIF2 $\alpha$ , along with other ER stress markers, is upregulated (24). Induction of ER stress is accompanied by hypersensitivity, and attenuation of ER stress by an inhibitor of soluble epoxide hydrolase (sEH) down-regulated ER stress markers, p-PERK, p-eIF2 $\alpha$ , and reversed mechanical hypersensitivity (24). Despite these intriguing observations, a direct link between  $eIF2\alpha$ phosphorylation and nociception has not been established. Using a transgenic mouse model with reduced phosphorylation (by  $\sim 50\%$ ) of eIF2 $\alpha$  (eIF2 $\alpha^{+/S51A}$ ), we show that p-eIF2 $\alpha$  controls thermal, but not mechanical, sensitivity via the regulation of transient receptor potential vanilloid receptor 1 (TRPV1) activity. Moreover, we find that eIF2a phosphorylation is induced in primary nociceptors in a chronic inflammation model and that it contributes to inflammatory pain hypersensitivity.

## Results

**p-eIF2** $\alpha$  **Is Increased in Dorsal Root Ganglia After Chronic Inflammation.** First, we examined the distribution of eIF2 $\alpha$  and its phosphorylated form, p-eIF2 $\alpha$ , in dorsal root ganglia (DRGs) and spinal cord. Immunostaining revealed neuronal expression of eIF2 $\alpha$ and p-eIF2 $\alpha$  in peptidergic (CGRP-positive), nonpeptidergic (IB4positive), TRPV1-positive small diameter, and NF200-positive large diameter neuronal cell bodies in DRGs (Fig. 1 *A* and *B*). Colocalization analysis showed that 19.7% of p-eIF2 $\alpha$ -positive cells express CGRP, 36.6% express IB4, 20.8% express TRPV1, and 47% express NF200 (Fig. 1*B*, *Bottom*). In the dorsal horn of the spinal cord, eIF2 $\alpha$  and p-eIF2 $\alpha$  were found in neurons, as they colocalized with the neuronal marker NeuN, but not with the astrocyte marker, glial fibrillary acidic protein (GFAP) (Fig. S1). However, in the spinal cord p-eIF2 $\alpha$  signal was rather weak and detected only in a small fraction of NeuN-positive neurons (Fig. S1).

To determine whether phosphorylation of eIF2 $\alpha$  is affected by chronic inflammation, we injected complete Freund's adjuvant (CFA) s.c. into the mouse hind paw (intraplantar injection) and measured the levels of p-eIF2 $\alpha$  in lumbar DRGs and dorsal horn of the spinal cord. Levels of p-eIF2 $\alpha$  were increased in DRGs, but not in the spinal cord, 1 d after the onset of inflammation, decreased subsequently, and returned to normal after 10 d (Fig. 1*C*). The alterations in p-eIF2 $\alpha$  concurred with the inflammationinduced changes in thermal and mechanical thresholds (Fig. 1*D*), raising the possibility that the increase in eIF2 $\alpha$  phosphorylation mediates the inflammatory hypersensitivity.

P-elF2 $\alpha$  Controls Thermal Sensitivity. Having established that p-elF2 $\alpha$ is increased in DRGs in response to chronic inflammation, we investigated its role in nociception. To this end, we used a transgenic knock-in (KI) mouse model (25), in which serine-51 is mutated to a nonphosphorylatable alanine residue in one allele  $(eIF2\alpha^{+/S51A})$ , homozygous KI mice are not viable), leading to a ~50% reduction in basal eIF2 $\alpha$  phosphorylation (Fig. 2A). Mechanical sensitivity in the von Frey and tail clip tests did not differ between  $eIF2\alpha^{+/S51A}$  mice and their WT littermates (Fig. 2B). However, thermal withdrawal and nocifensive behavior latencies were significantly prolonged in  $eIF2\alpha^{+/S51A}$  mice compared with WT mice in the radiant heat paw withdrawal, hot water tail withdrawal, and hot-plate tests ( $40.2 \pm 9.7\%$ ,  $44.0 \pm 13.1\%$ , and  $21.6 \pm$ 6.6% increase, respectively; Fig. 2C), indicating reduced sensitivity to noxious heat in  $eIF2\alpha^{+/S51A}$  mice. No difference in sensitivity to noxious cold was observed between WT and  $eIF2\alpha^{+/S51A}$  mice (Fig. 2D).  $eIF2\alpha^{+/S51A}$  mice also exhibited reduced inflammatory pain in the formalin test. Nocifensive (licking/shaking) behavior was significantly reduced (by  $31.9 \pm 6.0\%$ ) in  $eIF2\alpha^{+/S51A}$  mice during the late/tonic phase (10-60 min after formalin injection), compared



**Fig. 1.** eIF2 $\alpha$  is expressed in DRG neurons and its phosphorylation is increased in an inflammatory pain model. The distribution of total and p-eIF2 $\alpha$  in mouse lumbar DRG was examined using immunostaining. Total (*A*) and p-eIF2 $\alpha$  (B) were costained with CGRP, IB4, TRPV1, and NF200. Percent of p-eIF2 $\alpha$ -positive neurons expressing the markers is shown in *B* (*Bottom*). (C) Mice were injected (intraplantar) with CFA, and levels of eIF2 $\alpha$  phosphorylation were measured in DRGs at different time points after injection using Western blot analysis (n = 4 mice per condition). (*D*) CFA induces thermal (*Left*) and mechanical (*Right*) hypersensitivity as assessed in radiant heat paw withdrawal and von Frey assays, respectively (n = 5 males and 4 females per assay). Data are presented as mean  $\pm$  SEM. \*P < 0.05, \*\*P < 0.01 by Bonferroni post hoc test following one-way ANOVA. (Scale bar, 100 µm.) For distribution of eIF2 $\alpha$  and p-eIF2 $\alpha$  in the spinal cord, see Fig. S1.

with WT mice, whereas no differences were found in the early/ acute phase (0–10 min) between these groups (Fig. 2*E*). The behavioral differences occurred despite equal degrees of paw edema in the two genotypes (Fig. 2*E*). Taken together, these results demonstrate that p-eIF2 $\alpha$  is up-regulated in DRGs in response to chronic inflammation, and mice with reduced eIF2 $\alpha$  phosphorylation exhibit decreased heat sensitivity but not mechanical sensitivity.

As eIF2 $\alpha$  is phosphorylated by four different kinases (Fig. 3*A*), it was pertinent to study the effect of each kinase on p-eIF2 $\alpha$  and thermal threshold. Because HRI expression is very low in the nervous system (12), we examined sensitivity to noxious heat in *Perk*<sup>+/-</sup> (*Perk*<sup>-/-</sup> exhibits severe postnatal growth retardation) (26), *Pkr*<sup>-/-</sup>, and *Gcn*2<sup>-/-</sup> mice. *Perk* heterozygous had reduced p-eIF2 $\alpha$  in DRGs and decreased noxious heat sensation (43.0 ±7.7% increase in latency to withdrawal in the radiant heat paw withdrawal test; Fig. 3*B*). Mechanical thresholds in the von Frey test were not altered in *Perk*<sup>+/-</sup> mice, similar to *eIF2\alpha*<sup>+/S51A</sup> mice. *Gcn*2<sup>-/-</sup> and *Pkr*<sup>-/-</sup> mice did not display a significant reduction in p-eIF2 $\alpha$  level and sensitivity to noxious heat (Fig. 3 *C* and *D*); however, double KO mice for *Gcn*2 and *Pkr* (*Gcn*2/*Pkr* DKO) exhibited reduced p-eIF2 $\alpha$  levels (Fig. 3*C*) and elevated thermal thresholds (Fig. 3*E*). This finding suggests a redundant role for these two kinases.



**Fig. 2.** Noxious heat sensation is reduced in  $eIF2a^{+/551A}$  mice. (A) eIF2a phosphorylation is reduced in DRGs and spinal cord of  $eIF2a^{+/551A}$  mice.  $eIF2a^{+/551A}$  mice demonstrate no alterations in mechanical sensitivity (B; n = 4 males and 4 females per genotype, P > 0.05), whereas noxious heat sensation is significantly attenuated (C; n = 4 males and 4 females per genotype per assay, P < 0.05). (D) Sensitivity to cold is not changed in  $eIF2a^{+/551A}$  mice (n = 4 males and 6 females per genotype). (E) Nocifensive (licking/shaking) behavior is significantly reduced in formalin test during the late/tonic phase (10–60 min after formalin injection, P < 0.05), whereas no differences are found in the early/acute phase (n = 4 males and 4 females per genotype). Changes in paw weight, indicative of formalin-induced inflammation, are not different in  $eIF2a^{+/551A}$  mice (*E*, *Right*; P > 0.05). Data are presented as mean  $\pm$  SEM. \*P < 0.05, \*\*P < 0.01; ns, not significant by Student *t* test.

Next, we examined whether modulation of  $eIF2\alpha$  phosphorylation by drugs alters the thermal threshold.  $eIF2\alpha$  phosphorylation was decreased by an inhibitor of eIF2 $\alpha$  kinase, PKR (PKRi) (27). i.p. administration of PKRi over 3 d reduced noxious heat sensation in a dose-dependent manner, as indicated by the increased withdrawal latency in the radiant heat paw withdrawal test (Fig. 3F), with no effect on mechanical threshold. Conversely, when  $eIF2\alpha$  phosphorylation was increased by i.p. administration of Sal003, an inhibitor of the eIF2 $\alpha$  phosphatase complex, GADD34/PP1 (growth arrest and DNA-damageinducible 34/protein phosphatase1) (28) (Fig. 3A), thermal thresholds were decreased (Fig. 3 G and H), whereas mechanical thresholds were not affected. In summary, using genetic and pharmacological approaches, we show that decreasing  $eIF2\alpha$ phosphorylation reduces noxious heat sensation, whereas increasing p-eIF2 $\alpha$  levels engenders the opposite effect.

**TRPV1 Activity Mediates the Effect of Reduced p-elF2\alpha on Thermal Thresholds.** The strikingly specific impact of elF2 $\alpha$  phosphorylation on noxious heat sensation suggests that mechanisms controlling heat transduction might be selectively controlled. TRPV1 channels transduce noxious heat and are also implicated in inflammation-induced thermal hypersensitivity (29). TRPV1 activity is tightly regulated via gene expression and posttranslational mechanisms (30). We examined the possibility that TRPV1 mediates the effect of elF2 $\alpha$  phosphorylation on heat sensation by studying the impact of PKRi and Sal003 on thermal thresholds in  $Trpv1^{-/-}$  mice (29). PKRi increased thermal threshold in WT mice, but not in  $Trpv1^{-/-}$  mice (Fig. 4A). Conversely, Sal003 decreased

thermal threshold in WT mice, but not in  $TrpvI^{-/-}$  mice (Fig. 4*B*). These data demonstrate that eIF2 $\alpha$  phosphorylation controls thermal threshold in a TRPV1-dependent manner. To assess TRPV1 activity, we recorded TRPV1-dependent currents in sensory neurons from  $eIF2\alpha^{+/S51A}$  and WT mice. Capsaicin, a specific TRPV1 agonist, elicited significantly smaller currents in dissociated DRG neurons prepared from  $eIF2\alpha^{+/S51A}$  compared with WT mice (92% decrease in  $eIF2\alpha^{+/S51A}$  neurons; Fig. 4*C*). For whole-cell recordings, small-diameter (<30 µm) neurons were selected, and only capsaicin-sensitive neurons (~30% of all tested neurons in WT and  $eIF2\alpha^{+/S51A}$  groups) were included in the analysis. Resting membrane potential ( $V_{rest}$ ), input resistance ( $R_{in}$ ), and membrane capacitance ( $C_m$ ) were not different between WT and  $eIF2\alpha^{+/S51A}$  neurons (WT  $V_{rest}$  -49.74 ± 3.41 mV,  $eIF2\alpha^{+/S51A}$   $V_{rest}$  -46.79 ± 3.99 mV, P = 0.584; WT  $R_{in}$  686.21 ± 117.41 M $\Omega$ ,  $eIF2\alpha^{+/S51A}$   $R_{in}$  517.13 ± 38.41 M $\Omega$ , P = 0.193; WT



Fig. 3. eIF2 $\alpha$  kinases control thermal threshold. (A) eIF2 $\alpha$  kinases and phosphatase GADD34/PP1 (growth arrest and DNA-damage-inducible 34/ protein phosphatase1). (B)  $Perk^{+/-}$  mice show a decrease in p-elF2 $\alpha$  (n = 4 mice per genotype, P < 0.05) and in noxious heat sensitivity (n = 4 males and 4 females per genotype, P < 0.001). Gcn2<sup>-/-</sup> and Pkr<sup>-/-</sup> mice show no alteration in thermal latencies (C and D, respectively; n = 4 males and 4 females per genotype, P > 0.05). (E)  $Gcn2^{-/-} Pkr^{-/-}$  double KO mice show reduced noxious heat sensitivity (n = 3 males and 4 females per genotype, P < 0.05). eIF2 $\alpha$ phosphorylation in DRGs of Gcn2-/-, Pkr-/-, and Gcn2-/- Pkr-/- double KO mice is shown in C (n = 3 mice per genotype). PKR inhibitor (PKRi) (F) and Sal003 (G) were injected i.p. for 3 d daily, and thermal and mechanical thresholds were measured 30 min after the last injection (n = 4 males and 4 females per condition). (H) eIF2α phosphorylation in DRGs of mice injected with PKRi (1 mg/kg) and Sal003 (1 mg/kg) is shown (n = 5 mice per drug). Data are presented as mean ± SEM. \*P < 0.05, \*\*P < 0.01, \*\*\*P < 0.001; ns, not significant by Student t test and Bonferroni post hoc test following one-way ANOVA.



Fig. 4. eIF2α phosphorylation regulates thermal threshold via TRPV1. Modulation of eIF2 $\alpha$  phosphorylation in *Trpv1<sup>-/-</sup>* mice does not alter heat sensation. PKRi (1 mg/kg for 3 d daily, i.p.) elevates thermal threshold in WT but not in Trpv1<sup>-/-</sup> mice (A; n = 4 males and 4 females per genotype-drug condition). Sal003 (1 mg/kg for 3 d daily, i.p.) decreases thermal threshold in WT but not in Trpv1<sup>-/-</sup> mice (B, n = 4 males and 4 females per genotype-drug condition). For the radiant heat paw withdrawal test, the light beam was set to 20% of the maximal intensity in A and to 15% in B. Capsaicin (1  $\mu$ M) evokes smaller currents (C, n = 12 cells for  $eIF2\alpha^{+/S51A}$  and 10 for WT, from three different neuronal cultures per genotype) and smaller calcium tran-sients (D, n = 72 cells for eIF2 $\alpha$  <sup>+/S51A</sup> and n = 46 cells for WT, from four different neuronal cultures per genotype, using Fura-2 340/380-nm ratio) in cultured DRG neurons prepared from  $eIF2\alpha^{+/551A}$  compared with WT mice. Capsaicin (2.5  $\mu$ g), injected s.c. into the plantar surface of the hindpaw, elicited less nocifensive behaviors in  $eIF2\alpha^{+/S51A}$  compared with WT mice (E; n = 4 males and 4 females per genotype or drug), whereas in mice injected with Sal003 (1 mg/kg for 3 d daily, i.p.) capsaicin-induced pain behavior is increased (F). Western blot analysis shows that the TRPV1 protein levels are not altered in membrane and cytosolic fractions, as well as in total lysates of DRGs of  $eIF2a^{+/S51A}$  mice (G; n = 4 mice and genotype). TRPV1 surface levels were measured in DRG cultured neurons prepared from  $elF2a^{+/S51A}$  and WT mice using surface biotinylation assay (H; n = 3/group). Data are presented as mean  $\pm$  SEM. \*P < 0.05, \*\*P < 0.01, \*\*\*P < 0.001 by Student t test and Student t test following two-way (genotype  $\times$  drug) ANOVA.

 $C_m$  13.67 ± 2.18 pF,  $eIF2\alpha^{+/S51A}$   $C_m$  14.68 ± 1.95 pF, P = 0.735, n = 8/genotype). Moreover, using calcium imaging, we observed smaller capsaicin-induced calcium transients in cultured  $eIF2\alpha^{+/S51A}$  DRG neurons compared with WT neurons (Fig. 4D). The cell body diameter of the responding neurons was not different between the two genotypes (WT 19.18 ± 0.45 µm, n =53,  $eIF2\alpha^{+/S51A}$  19.32 ± 0.39 µm, n = 88, P = 0.81). Consistent with these results, intraplantar s.c. administration of capsaicin induced significantly less nocifensive behavior in  $eIF2\alpha^{+/S51A}$ compared with WT mice (Fig. 4E). Conversely, mice with high p-eIF2 $\alpha$  levels, following Sal003 injections, exhibited increased nociceptive responses to capsaicin (Fig. 4F). Despite the reduction in the amplitude of TRPV1-mediated currents in  $eIF2a^{+/S51A}$ neurons, Western blot analysis showed that protein levels of TRPV1 in cytosolic and membrane fractions of DRG lysates from  $eIF2a^{+/S51A}$  mice were not changed compared with WT mice (Fig. 4G). To examine whether trafficking of TRPV1 to the cell surface is affected by eIF2a phosphorylation, we used a surface biotinylation assay followed by Western blot analysis of TRPV1. We found no differences in TRPV1 amounts on the cell surface (Fig. 4H), indicating that TRPV1 activity, but not protein levels or trafficking to the plasma membrane, is reduced in  $eIF2a^{+/S51A}$  neurons. Taken together, these data indicate that TRPV1 is an important mediator of the effect of p-eIF2a on the thermal threshold and suggest that TRPV1 activity is modulated by eIF2a phosphorylation.

**PKR-Mediated elF2** $\alpha$  **Phosphorylation Contributes to Thermal Hypersensitivity.** After establishing that eIF2 $\alpha$  phosphorylation regulates thermal sensation, we studied whether CFA-induced increase in p-eIF2 $\alpha$  contributes to thermal hypersensitivity. eIF2 $\alpha$  kinase, PKR, has been implicated in inflammatory responses (31). Thus, we assessed PKR activation in DRGs following CFA injection and found that levels of p-PKR were significantly increased (Fig. 54), raising the possibility that the increase in p-eIF2 $\alpha$  is mediated via PKR activation. Consequently, we assessed thermal and mechanical hypersensitivity of *Pkr<sup>-/-</sup>* mice after CFA-induced inflammation. *Pkr<sup>-/-</sup>* mice exhibited reduced thermal,



Fig. 5. TNF- $\alpha$ -mediated PKR activation elevates p-elF2 $\alpha$ . (A) Phosphorylation of PKR (Thr451) was measured at different time points after CFA injection (n = 4 mice). (B) CFA-induced thermal hyperalgesia is attenuated in Pkr<sup>-/-</sup> mice compared with WT mice, whereas mechanical hyperalgesia is not changed (n = 4 males and 4 females per genotype). (C) P-eIF2 $\alpha$  in DRGs was measured at different time points after CFA injection in WT and Pkr<sup>-/-</sup> mice. (D) HEK293 cells were treated with TNF- $\alpha$  (100 ng/mL) for 15, 30, and 60 min, and p-eIF2 $\alpha$  was measured (n = 3, P < 0.05). (E) TNF- $\alpha$  (20 ng in 5  $\mu$ L PBS + 0.5% BSA) was injected i.t. three times every 3 h in WT and Pkr<sup>-/-</sup> mice, and the p-elE2 $\alpha$  was measured in lumbar DRGs 30 min after the last injection TNF- $\alpha$  induced an increase in p-eIF2 $\alpha$  in WT but not in  $Pkr^{-/-}$  mice (n = 5 females per genotype per drug). (F) TNF- $\alpha$  (20 ng, i.t., three injections at 3-h intervals) elicited bigger thermal hyperalgesia in WT mice compared with  $Pkr^{-/-}$  mice (n = 4 males and 4 females per genotype per drug, P < 0.05). Data are presented as mean  $\pm$  SEM. \*P < 0.05, \*\*P < 0.01, \*\*\*P < 0.001 by Bonferroni post hoc test following one-way ANOVA and Student t test following two-way (genotype  $\times$  drug) ANOVA.

but not mechanical, pain hypersensitivity after CFA injection (Fig. 5*B*). Moreover, p-eIF2 $\alpha$  increased after CFA injection in DRGs of WT mice, but not in *Pkr<sup>-/-</sup>* mice (Fig. 5*C*). These data demonstrate that PKR is required for the up-regulation of p-eIF2 $\alpha$  following CFA and contributes to thermal hyperalgesia.

TNF- $\alpha$  induces a robust up-regulation of eIF2 $\alpha$  phosphorylation in cultured cells and in the nervous system via activation of PKR (32–36). Because TNF- $\alpha$  is a major proinflammatory cytokine (37-40), which plays a critical role in the pathogenesis of inflammatory pain (41, 42), we examined whether inflammationinduced TNF- $\alpha$  promotes eIF2 $\alpha$  phosphorylation. First, we showed that TNF- $\alpha$  elevates p-eIF2 $\alpha$  in HEK293 cells, replicating previous studies (Fig. 5D). Importantly, intrathecally delivered TNF- $\alpha$  increased p-eIF2 $\alpha$  in DRGs of WT mice, but not  $Pkr^{-/-}$  mice (Fig. 5E), supporting the idea that TNF- $\alpha$  stimulates p-eIF2a via PKR. In accordance with previous reports, TNF-a (i.t.) induced heat hyperalgesia in WT mice (37), whereas in mice this hyperalgesia was significantly attenuated (Fig.  $Pkr^{-/-}$ 5F). Taken together, these results demonstrate that TNF- $\alpha$ - and PKR-dependent eIF2α phosphorylation contributes to chronic inflammation-induced thermal hypersensitivity.

## Discussion

Here we describe a previously unrecognized role for the cellular stress response pathway in nociception. Transgenic mice with decreased eIF2 $\alpha$  phosphorylation (eIF2 $\alpha^{+/S51A}$ ) exhibited reduced responses to noxious heat and attenuated nocifensive behavior in the late phase of formalin test, whereas cold sensitivity and mechanical thresholds were not altered. The noxious heat-specific phenotype was recapitulated in transgenic mice in which the eIF2a kinases PERK and PKR/GCN2 were reduced or knocked out, as well as in response to pharmacological manipulation of  $eIF2\alpha$  phosphorylation. Reducing  $eIF2\alpha$  phosphorylation with a PKR inhibitor attenuated noxious heat sensitivity, whereas increasing eIF2a phosphorylation with Sal003 had the opposite effect. Our findings indicate that the effect of p-eIF2 $\alpha$ on thermal nociception is mediated via modulation of TRPV1 activity. First, we show that pharmacological modulation of eIF2a phosphorylation altered noxious heat sensation in WT mice, but had no effect in mice lacking TRPV1. Second, capsaicin-induced TRPV1-mediated currents and pain behavior were greatly reduced in  $eIF2\alpha^{+/S51A}$  mice. Taken together, these data demonstrate that the eIF2 $\alpha$  pathway controls noxious heat sensation via TRPV1. Because we found no evidence of alterations in TRPV1 protein levels or trafficking to the membrane, we postulate that the mechanism by which eIF2a phosphorylation affects thermal sensation involves modulation of TRPV1 activity. Under inflammatory conditions, TRPV1 can be sensitized by numerous inflammatory mediators (e.g., bradykinin, nerve growth factor, prostaglandins, serotonin, and histamine), via multiple agonists and modulators [protein kinase A (PKA), protein kinase C (PKC), metabolites of the cyclooxygenase (COX), lipoxygenase (LOX), and cytochrome-P450 (CYP)-pathways, phospholipids, protons, and heat, among others], leading to the reduction in the activation threshold and eventually to pain hypersensitivity (43, 44). The modulators, which mediate the sensitization of TRPV1 activity by p-eIF2a, remain to be determined. eIF2a phosphorylation promotes translation of ATF4 mRNA (9, 11). Interestingly, ATF4 is increased in DRGs following facet joint distraction (45). Moreover, ATF4 transcriptionally activates ATF3 expression, which is a well-known cellular marker of nerve injury (46). This evidence suggests that some of the effects of p-eIF2 $\alpha$  on thermal thresholds could be mediated by ATF4/ATF3 axis.

We show that TNF- $\alpha$  induces p-eIF2 $\alpha$  in WT but not in *Pkr<sup>-/-</sup>*mice, indicating that TNF- $\alpha$  up-regulates p-eIF2 $\alpha$  via PKR. PKR is activated after CFA injection, suggesting that TNF- $\alpha$ -mediated activation of PKR contributes to p-eIF2 $\alpha$  up-regulation and thermal hypersensitivity. Our results do not exclude the involvement of other eIF2 $\alpha$  kinases. For example, ER stress induces a robust PERK activation (19), which has a strong effect on eIF2 $\alpha$  phosphorylation and thermal sensitivity (Fig. 3*B*).

A recent study found that hyperglycemia, activation of unfolded protein response (UPR), or dysregulation of calcium homeostasis induce ER stress in primary sensory neurons, as evident by the activation of PERK (and eIF2 $\alpha$ ), inositol-requiring enzyme 1a (IRE1a), ATF6, MAPK (p38 and JNK), and autophagy (LC3) (24). Whether the ER stress-induced mechanical pain is caused by elevated p-eIF2 $\alpha$  or through other mechanisms was not documented. Because elevated p-eIF2 $\alpha$  affects thermal, but not mechanical thresholds, it seems unlikely that the effects of ER stress on nociception are mediated via p-eIF2 $\alpha$ , but could be attributed to the activation of p38 and JNK or to other ER stress-dependent mechanisms. Increased p-eIF2 $\alpha$  was also documented in the sciatic nerve of rats with presumed neuropathic pain (47); however, the impact of this phosphorylation event on nociception has not been investigated.

Recent preclinical studies concluded that modulators of eIF2 $\alpha$  phosphorylation might have therapeutic potential in treatment of several cellular stress-related pathologies such as Alzheimer's disease (22, 48), prion diseases (23), diabetes (49), Huntington's disease (50), and amyotrophic lateral sclerosis (51). It will be important to consider the effect of eIF2 $\alpha$  phosphorylation on thermal nociception while developing clinically applicable compounds to the latter maladies.

In summary, we uncovered a previously unknown role for the cellular stress response pathway in nociception. This knowledge can be used to develop strategies to treat conditions associated with altered heat sensation, most notably burn pain, and should be considered while introducing  $eIF2\alpha$  modulators to clinical practice.

## **Experimental Procedures**

**Behavioral Experiments, Electrophysiological Recordings, and Calcium Imaging.** See *SI Experimental Procedures* for details of the experimental procedures. All procedures complied with Canadian Council on Animal Care guidelines and were approved by McGill University's Downtown Animal Care Committee.

**Drugs.** PKR inhibitor (PKRi) and Sal003 were purchased from Calbiochem and dissolved in 30% polyethylene glycol in saline. Capsaicin was purchased from Sigma and dissolved in ethanol. TNF- $\alpha$  was purchased from Kamiya Biomedical.

Western Blotting and Immunohistochemistry. Proteins were resolved on SDSpolyacrylamide gels using standard techniques. See *SI Experimental Procedures* for details of the experimental procedures and antibodies used.

**Statistical Analyses.** All results are expressed as mean  $\pm$  SEM. All statistical comparisons were made with either the Student *t* test or a one-way ANOVA followed by between-group comparisons using the Bonferroni post hoc test, unless otherwise indicated, with *P* < 0.05 as the significance criterion. Power analyses were not possible because we had no a priori expectation of effect size but rather were informed by normative practices in the pain field (52).

ACKNOWLEDGMENTS. We thank A. Sylvestre, S. Perreault, C. Lister, and I. Harvey for technical assistance and I. Daou for technical advice. This work was supported by Canadian Institutes of Health Research (CIHR) Operating Grant MOP-114994 (to N.S.), an unrestricted gift from the Louise and Alan Edwards Foundation (to J.S.M.), CIHR Grant FDN-143337 (to C.W.B.), and the James McGill Chair Program (C.W.B.).

- Hinnebusch AG, Ivanov IP, Sonenberg N (2016) Translational control by 5'-untranslated regions of eukaryotic mRNAs. Science 352(6292):1413–1416.
- Sonenberg N, Hinnebusch AG (2009) Regulation of translation initiation in eukaryotes: Mechanisms and biological targets. Cell 136(4):731–745.

Downloaded from https://www.pnas.org by UNIVERSITY OF TORONTO on July 2, 2022 from IP address 142.150.190.39.

Holcik M, Sonenberg N (2005) Translational control in stress and apoptosis. Nat Rev Mol Cell Biol 6(4):318–327.

<sup>2.</sup> Ron D (2002) Translational control in the endoplasmic reticulum stress response. J Clin Invest 110(10):1383–1388.

- Donnelly N, Gorman AM, Gupta S, Samali A (2013) The elF2alpha kinases: Their structures and functions. *Cell Mol Life Sci* 70(19):3493–3511.
- Raven JF, Koromilas AE (2008) PERK and PKR: Old kinases learn new tricks. *Cell Cycle* 7(9):1146–1150.
- Wek RC, Jiang HY, Anthony TG (2006) Coping with stress: eIF2 kinases and translational control. *Biochem Soc Trans* 34(Pt 1):7–11.
- Trinh MA, Klann E (2013) Translational control by eIF2α kinases in long-lasting synaptic plasticity and long-term memory. *Neurobiol Learn Mem* 105:93–99.
- Harding HP, et al. (2000) Regulated translation initiation controls stress-induced gene expression in mammalian cells. Mol Cell 6(5):1099–1108.
- Han J, et al. (2013) ER-stress-induced transcriptional regulation increases protein synthesis leading to cell death. Nat Cell Biol 15(5):481–490.
- 11. Rutkowski DT, Kaufman RJ (2003) All roads lead to ATF4. Dev Cell 4(4):442-444.
- Costa-Mattioli M, Sossin WS, Klann E, Sonenberg N (2009) Translational control of long-lasting synaptic plasticity and memory. *Neuron* 61(1):10–26.
- Trinh MA, et al. (2014) The eIF2α kinase PERK limits the expression of hippocampal metabotropic glutamate receptor-dependent long-term depression. *Learn Mem* 21(5):298–304.
- Di Prisco GV, et al. (2014) Translational control of mGluR-dependent long-term depression and object-place learning by eIF2α. Nat Neurosci 17(8):1073–1082.
- Hetz C, Mollereau B (2014) Disturbance of endoplasmic reticulum proteostasis in neurodegenerative diseases. Nat Rev Neurosci 15(4):233–249.
- Ozcan U, et al. (2004) Endoplasmic reticulum stress links obesity, insulin action, and type 2 diabetes. Science 306(5695):457–461.
- Wang M, Kaufman RJ (2014) The impact of the endoplasmic reticulum protein-folding environment on cancer development. Nat Rev Cancer 14(9):581–597.
- Koromilas AE (2015) Roles of the translation initiation factor eIF2α serine 51 phosphorylation in cancer formation and treatment. *Biochim Biophys Acta* 1849(7): 871–880.
- Segev Y, Michaelson DM, Rosenblum K (2013) ApoE ε4 is associated with eIF2α phosphorylation and impaired learning in young mice. Neurobiol Aging 34(3): 863–872.
- Chang RC, Wong AK, Ng HK, Hugon J (2002) Phosphorylation of eukaryotic initiation factor-2alpha (elF2alpha) is associated with neuronal degeneration in Alzheimer's disease. *Neuroreport* 13(18):2429–2432.
- O'Connor T, et al. (2008) Phosphorylation of the translation initiation factor elF2alpha increases BACE1 levels and promotes amyloidogenesis. *Neuron* 60(6):988–1009.
- Ma T, et al. (2013) Suppression of eIF2α kinases alleviates Alzheimer's disease-related plasticity and memory deficits. Nat Neurosci 16(9):1299–1305.
- 23. Moreno JA, et al. (2012) Sustained translational repression by elF2 $\alpha$ -P mediates prion neurodegeneration. *Nature* 485(7399):507–511.
- Inceoglu B, et al. (2015) Endoplasmic reticulum stress in the peripheral nervous system is a significant driver of neuropathic pain. Proc Natl Acad Sci USA 112(29):9082–9087.
- Scheuner D, et al. (2001) Translational control is required for the unfolded protein response and in vivo glucose homeostasis. *Mol Cell* 7(6):1165–1176.
- 26. Zhang P, et al. (2002) The PERK eukaryotic initiation factor 2 alpha kinase is required for the development of the skeletal system, postnatal growth, and the function and viability of the pancreas. *Mol Cell Biol* 22(11):3864–3874.
- Jammi NV, Whitby LR, Beal PA (2003) Small molecule inhibitors of the RNA-dependent protein kinase. *Biochem Biophys Res Commun* 308(1):50–57.
- Novoa I, Zeng H, Harding HP, Ron D (2001) Feedback inhibition of the unfolded protein response by GADD34-mediated dephosphorylation of elF2alpha. J Cell Biol 153(5):1011–1022.
- 29. Caterina MJ, et al. (2000) Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science* 288(5464):306–313.
- Planells-Cases R, Valente P, Ferrer-Montiel A, Qin F, Szallasi A (2011) Complex regulation of TRPV1 and related thermo-TRPs: Implications for therapeutic intervention. *Adv Exp Med Biol* 704:491–515.
- 31. Kang R, Tang D (2012) PKR-dependent inflammatory signals. Sci Signal 5(247):pe47.
- Mantuano E, et al. (2011) The unfolded protein response is a major mechanism by which LRP1 regulates Schwann cell survival after injury. J Neurosci 31(38): 13376–13385.

- Sharma B, Altman JK, Goussetis DJ, Verma AK, Platanias LC (2011) Protein kinase R as mediator of the effects of interferon (IFN) gamma and tumor necrosis factor (TNF) alpha on normal and dysplastic hematopoiesis. J Biol Chem 286(31):27506–27514.
- Hu S, et al. (2007) Translational inhibition of colonic epithelial heat shock proteins by IFN-gamma and TNF-alpha in intestinal inflammation. *Gastroenterology* 133(6): 1893–1904.
- Gilbert SJ, Duance VC, Mason DJ (2004) Does protein kinase R mediate TNF-alpha- and ceramide-induced increases in expression and activation of matrix metalloproteinases in articular cartilage by a novel mechanism? *Arthritis Res Ther* 6(1):R46–R55.
- Lourenco MV, et al. (2013) TNF-α mediates PKR-dependent memory impairment and brain IRS-1 inhibition induced by Alzheimer's β-amyloid oligomers in mice and monkeys. *Cell Metab* 18(6):831–843.
- Zhang L, et al. (2011) TNF-α contributes to spinal cord synaptic plasticity and inflammatory pain: Distinct role of TNF receptor subtypes 1 and 2. Pain 152(2):419–427.
- Ohtori S, Takahashi K, Moriya H, Myers RR (2004) TNF-alpha and TNF-alpha receptor type 1 upregulation in glia and neurons after peripheral nerve injury: Studies in murine DRG and spinal cord. *Spine* 29(10):1082–1088.
- Schäfers M, Geis C, Svensson CI, Luo ZD, Sommer C (2003) Selective increase of tumour necrosis factor-alpha in injured and spared myelinated primary afferents after chronic constrictive injury of rat sciatic nerve. *Eur J Neurosci* 17(4):791–804.
- Uçeyler N, Tscharke A, Sommer C (2008) Early cytokine gene expression in mouse CNS after peripheral nerve lesion. *Neurosci Lett* 436(2):259–264.
- Xu ZZ, et al. (2010) Resolvins RvE1 and RvD1 attenuate inflammatory pain via central and peripheral actions. Nat Med 16(5):592–597.
- Cunha FQ, Poole S, Lorenzetti BB, Ferreira SH (1992) The pivotal role of tumour necrosis factor alpha in the development of inflammatory hyperalgesia. Br J Pharmacol 107(3):660–664.
- Morales-Lázaro SL, Simon SA, Rosenbaum T (2013) The role of endogenous molecules in modulating pain through transient receptor potential vanilloid 1 (TRPV1). J Physiol 591(13):3109–3121.
- Sisignano M, Bennett DL, Geisslinger G, Scholich K (2014) TRP-channels as key integrators of lipid pathways in nociceptive neurons. Prog Lipid Res 53:93–107.
- Dong L, Guarino BB, Jordan-Sciutto KL, Winkelstein BA (2011) Activating transcription factor 4, a mediator of the integrated stress response, is increased in the dorsal root ganglia following painful facet joint distraction. *Neuroscience* 193:377–386.
- Bráz JM, Basbaum AI (2010) Differential ATF3 expression in dorsal root ganglion neurons reveals the profile of primary afferents engaged by diverse noxious chemical stimuli. *Pain* 150(2):290–301.
- Melemedjian OK, et al. (2011) Targeting adenosine monophosphate-activated protein kinase (AMPK) in preclinical models reveals a potential mechanism for the treatment of neuropathic pain. *Mol Pain* 7:70.
- Fullwood MJ, Zhou W, Shenolikar S (2012) Targeting phosphorylation of eukaryotic initiation factor-2α to treat human disease. Prog Mol Biol Transl Sci 106:75–106.
- Back SH, et al. (2009) Translation attenuation through elF2alpha phosphorylation prevents oxidative stress and maintains the differentiated state in beta cells. *Cell Metab* 10(1):13–26.
- Reijonen S, Putkonen N, Nørremølle A, Lindholm D, Korhonen L (2008) Inhibition of endoplasmic reticulum stress counteracts neuronal cell death and protein aggregation caused by N-terminal mutant huntingtin proteins. *Exp Cell Res* 314(5):950–960.
- Saxena S, Cabuy E, Caroni P (2009) A role for motoneuron subtype-selective ER stress in disease manifestations of FALS mice. Nat Neurosci 12(5):627–636.
- Mogil JS, et al. (2006) Screening for pain phenotypes: Analysis of three congenic mouse strains on a battery of nine nociceptive assays. *Pain* 126(1-3):24–34.
- Abraham N, et al. (1999) Characterization of transgenic mice with targeted disruption of the catalytic domain of the double-stranded RNA-dependent protein kinase, PKR. J Biol Chem 274(9):5953–5962.
- Chaplan SR, Bach FW, Pogrel JW, Chung JM, Yaksh TL (1994) Quantitative assessment of tactile allodynia in the rat paw. J Neurosci Methods 53(1):55–63.
- Gong K, Kung LH, Magni G, Bhargava A, Jasmin L (2014) Increased response to glutamate in small diameter dorsal root ganglion neurons after sciatic nerve injury. *PLoS One* 9(4):e95491.
- Côté S, Ribeiro-da-Silva A, Cuello AC (1993) Current protocols for light microscopy immunocytochemistry. *Immunohistochemistry II*, ed Cuello AC (Wiley, Chichester, UK), pp 147–168.